

Pre-master Probability and Statistics (2DI50), 11 May 2007, 9.00-12.00

- At the exam it is allowed to use the 'Statistisch Compendium' and a pocket calculator.
- The exam has 8 exercises with 20 items.
- The answers have to be in ENGLISH. Give not just answers, but also arguments.
- For each item one get get two points. In total one can get 40 points.

Problem 1:

Two balls are extracted from an urn containing white and black balls. After the first extraction, the selected ball is replaced in the urn. The probability of extracting a white ball is thus constant at each trial and it will be denoted by p , with $0 < p < 1$. Define the following two discrete random variables:

$$X = \begin{cases} 1 & \text{if the two extracted balls have the same color} \\ 0 & \text{otherwise} \end{cases}$$

$$Y = (\text{number of extracted white balls}) - (\text{number of extracted black balls})$$

(a) Compute the joint probability mass function $f_{X,Y}(x, y) = \mathbb{P}(X = x, Y = y)$

$P(X = x, Y = y)$	Y	$P(X = x)$
	· · ·	
	· · · ·	·
X	· · · ·	·
	· · ·	·
$P(Y = y)$	· · ·	

- (b) Find the values of p for which X and Y are uncorrelated, namely the covariance is zero.
- (c) Decide wheter X and Y independent for the values of p found in the previous item.

Problem 2:

A machine contains five components whose lifetime (in hours) has exponential distribution with mean 1000. The producer of the machine is considering to offer a guarantee that no more than two of the original components will have to be replaced during the first 1000 hours of use. Find the probability that such a guarantee will be violated in the following two cases:

- (a) the components wear out independently and when a component does fail then the replacement used is of particularly high quality and will certainly last for the 1000 hours.
- (b) the components wear out independently and no assumptions are made on the replacement components.

Problem 3:

A printer receives, on average, 5 jobs per minutes. Assume the number of assigned printing jobs has a Poisson distribution. Find the probability:

- (a) that in a 1 minute period no jobs are received
- (b) that in a 2 minutes period fewer than 4 jobs are received
- (c) that in 20 minutes no more than 102 jobs are received
- (d) that out of five separate 1 minute periods there are exactly four in which 2 or more jobs are received.

Problem 4:

The concentration C of a certain active agent in a pesticide must not exceed 12 part per million. The pesticide is made up in batches and in each batch the concentration is normally distributed, with mean 8 part per million and standard deviation 1.5 part per million.

- (a) What proportion of batches exceeds the permitted maximum?

Two batches chosen at random and independently of each other are mixed so that the resulting concentration X is the average of the two concentration in the constituent batches: $X = \frac{1}{2}(C_1 + C_2)$.

- (b) What is the probability that such a mixed sample has concentration value between 6 and 12 parts per million?
- (c) What is the probability that the concentration of the mixed sample falls below 3 part per million (the level at which the pesticide cease to have any effect)?

Problem 5:

A person plays chess against a computer once a day. He can set the difficulty to 3 levels: easy, medium or difficult. The player sets an “easy” level with probability $1/2$, a “medium” level with probability $1/4$ and a “difficult” level with probability $1/4$. If he plays an easy level game he has an 80% chance of winning; corresponding figures for medium and difficult level are 40% and 60% respectively.

- (a) Find the probability that, on a given day, the player wins.
- (b) If, on a particular day, the player has lost, which level has he most likely played?

Problem 6:

Let $(X_1, X_2, X_3, X_4, X_5)$ be a random sample from a random variable X with expectation value $E(X) = \mu$ and variance $V(X) = \sigma^2$. Consider the following three estimators for the expected value μ

1. $T_1(X_1, X_2, X_3, X_4, X_5) = X_1$,
2. $T_2(X_1, X_2, X_3, X_4, X_5) = \frac{1}{3}(X_1 + X_3 + X_5)$,
3. $T_3(X_1, X_2, X_3, X_4, X_5) = \frac{1}{6}(X_1 + X_2 + X_3 + X_4 + X_5)$.

- (a) Are the three estimators unbiased? Compute the bias.
- (b) Compute the Mean Squared Error of the three estimators.
- (c) Suppose $|\mu| < 2$ and $\sigma^2 = 1$. Which is the estimator with the smallest MSE ?

Problem 7:

A pencil producer thinks that his pencil machine actually produces pencils of a mean length of $\mu = 17\text{cm}$. He assumes that the machine produces pencils with normally distributed lengths. The variance is unknown. He measures the lengths of five pencils from the production and obtains lengths of 19.2 cm, 17.4 cm, 18.5 cm, 16.5 cm, 18.9 cm.

- (a) Compute sample mean \bar{x} and sample variance s^2 .
- (b) Construct a 95 % - confidence interval for the mean pencil length. Does the result contradict the producer’s opinion ?

Problem 8:

Upgrading a certain software package requires installation of 68 new files. Files are installed consecutively. The installation time is random, but on the average, it takes 15 sec to install one file, with a variance of 11 sec²

- (a) What is the probability that the whole package is updated in less than 16 minutes?
- (b) A new version of the package is released. It requires only N new files to be installed, and it is premised that 95% of the time upgrading takes less than 10 minutes. Given this information, compute N .